

**TRAINING PROTOCOLS FOR THE EMERGENCY ADMINISTRATION  
OF EPINEPHRINE**

**Lucille E. Davy**  
*Commissioner of Education*

**Barbara Gantwerk**  
*Assistant Commissioner*  
*Division of Student Services*

**Susan Martz**  
*Director*  
*Office of Educational Support Services*

**NEW JERSEY DEPARTMENT OF EDUCATION**  
**P. O. BOX 500**  
**TRENTON, NEW JERSEY 08625-0500**  
**(609) 292-5935**

*September 2008*  
*PTM Number - 1506.57*

# STATE BOARD OF EDUCATION

JOSEPHINE E. HERNANDEZ ..... President	Union
ARCELIO APONTE..... Vice President	Middlesex
RONALD K. BUTCHER.....	Gloucester
KATHLEEN A. DIETZ .....	Somerset
DEBRA ECKERT-CASHA.....	Morris
EDITHE FULTON.....	Ocean
ERNEST P. LEPORE .....	Hudson
FLORENCE MCGINN.....	Hunterdon
KENNETH J. PARKER.....	Camden
DOROTHY STRICKLAND.....	Essex

Lucille E. Davy, Commissioner of Education  
Secretary, State Board of Education

# TABLE OF CONTENTS

## ACKNOWLEDGEMENTS

### **PART A**

---

I. INTRODUCTION	p. 1
II. TRAINING STANDARDS	p. 2
RECOGNIZING SYMPTOMS OF ANAPHYLAXIS	p. 2
III. STANDARDS AND PROCEDURES FOR THE EMERGENCY USE OF AUTO-INJECTORS	p. 4
IV. EMERGENCY FOLLOW-UP PROCEDURES	p. 7
V. POLICIES AND PROCEDURES	p. 7

### **PART B**

---

I. THE ROLE OF SCHOOL BOARDS AND ADMINISTRATORS	p. 9
II. THE ROLES OF THE SCHOOL NURSE	p. 10
III. THE ROLE OF THE DELEGATE	p. 12

### **PART C**

---

APPENDIX	P. 13
----------	-------

## ACKNOWLEDGEMENTS

The New Jersey Department of Education would like to acknowledge the collaborative effort that resulted in the development of this document and wishes to thank the members of the School Health Services Committee for their contributions, without which this project would not have been possible. The department also acknowledges the following staff for their work in preparing and developing this document: LaCoyya Weathington, Office of Educational Support Services Division of Student Services, Sarah Kleinman, Office of Educational Support Services Division of Student Services, and Elaine Lerner, Office of Special Education Programs Division of Student Services, and Susan Martz, Office of Educational Support Services Division of Student Services.

Special thanks to the New Jersey Board of Nursing and George Herbert, Executive Director, for its review of this document.

### Department of Education

Elaine Lerner  
Chrys Harttraft  
Beverly Hetrick  
Sarah Kleinman  
Heather Mills-Pevonis  
Daryl Minus-Vincent  
LaCoyya Weathington

### Department of Health and Senior Services

Cynthia Collins  
Janet DeGraaf  
Antonia Farrell  
Judith Hall  
Eric Hicken  
Linda Jones-Hicks  
Nancy Kelly Goodstein  
Kathleen Lutz  
Jenish Sudhakaran

### Department of Human Services

Margaret M Bennett  
Cindy Rogers  
Carol Siminski

### Certified School Nurses

Margaret Bush  
Susan Cohen  
Margaret Dooley (school nurse)  
Nan Masterson  
Carole Paladino

Beverly Stern

### NJ State Nurses Association

Andrea Aughenbaugh  
Carolyn Torre

### School Physicians

Jack Kripsak, MD  
Steven Rice, MD  
Wayne Yankus, MD

### Non-Public School Representatives

Deborah Cornell  
Dr. George V. Corwell  
Mary-Ann Gugliemella  
JoAnn Tier  
Susan Vincent

### Food Allergy and Anaphylaxis Assoc. of NJ

Barbara Cullori  
Allison Inerro  
Robert Pacenza  
Scott Sicherer

### NJ Education Association

Wayne Dibofsky

### NJ Principals and Supervisors Association

Deborah Bradley



## **PART A: TRAINING STANDARDS FOR THE ADMINISTRATION OF EPINEPHRINE VIA AUTO-INJECTORS**

*This document provides model training standards for the administration of epinephrine via auto-injectors in accordance with New Jersey P.L. 2007, c. 57. These standards are intended to provide guidelines for training school personnel who have volunteered to serve as school nurse designees and receive such training.*

### **I. INTRODUCTION**

**In 2007, the New Jersey legislature amended N.J.S.A. 18A:40-12.3-12.6 adding provisions that address the administration of epinephrine to students in New Jersey schools. The law requires the following:**

1. Trained designees for students who may require the emergency administration of epinephrine by auto-injector for anaphylaxis when the school nurse is not available;
2. Secure but unlocked storage of epinephrine in locations easily accessible by the school nurse and designee(s), to ensure prompt availability in the event of an allergic emergency at school or at a school-sponsored function (A school-sponsored function means any activity, event or program occurring on or off school grounds, whether during or outside of regular school hours, that is organized and/or supported by the school as per N.J.A.C. 6A:16-1.3);
3. Availability of the school nurse or designee(s) on site at the school and at school-sponsored functions in the event of an allergic reaction;
4. Transportation of the student to a hospital emergency room by emergency services personnel after the administration of epinephrine;
5. The establishment and dissemination of guidelines for the development of a policy by a school district or nonpublic school for the management of food allergies in the school setting and the emergency administration of epinephrine to students; and
6. Training protocols to assist the school nurse in recruiting and training additional school employees as volunteer designees.

**NOTE: N.J.A.C. 6A:16-1.4(a)1, 2 and 7 require school districts to develop policies and procedures for the care of any student who becomes injured or ill while at school or during participation in school-sponsored activities. These policies must delineate provisions for transportation and the emergency administration of epinephrine.**

## II. TRAINING STANDARDS

The school nurse is responsible for delegating the administration of epinephrine to school personnel in the event of an emergency. As such, the nurse must assess the competency of personnel to whom a task has been delegated. To do so, the school nurse must provide training for all school personnel responsible for the administration of epinephrine via an auto-injector in an emergency. This training must be provided at least annually and must include the information provided in this document.

<i>Common causes of anaphylaxis include:</i>	<i>Less common causes of anaphylaxis include:</i>
<ul style="list-style-type: none"> <li>✦ Food (most commonly - peanuts, walnuts, pecans fish, shellfish, eggs, milk, soy, and wheat)</li> <li>✦ Medication</li> <li>✦ Insect stings</li> <li>✦ Latex</li> </ul>	<ul style="list-style-type: none"> <li>✦ Exercise</li> <li>✦ Food-dependent exercise-induced anaphylaxis (occurs when a person eats a specific food and exercises within three to four hours after eating)</li> <li>✦ Idiopathic anaphylaxis (anaphylaxis with no apparent cause)</li> </ul>

### RECOGNIZING SYMPTOMS OF ANAPHYLAXIS.

When a school nurse has recruited a delegate to administer epinephrine in an emergency, the nurse must then provide training to prepare the delegate for this responsibility. Part of the training should include a review of the student's Individualized Emergency Healthcare Plan (IEHP) so that the delegate will be aware of the symptoms that may trigger an allergic reaction in that specific student. The delegate must become familiar with recognizing the symptoms of allergic or anaphylactic reactions and triggers that cause these reactions for each student as delineated in the medical orders of the student's medical home.

#### Examples of symptoms that may occur during an allergic reaction/anaphylaxis

(The severity of symptoms can change quickly.)

Mouth: Itching, tingling, or swelling of lips, tongue, and mouth  
 Skin: Hives, itchy rash, swelling on the face or extremities  
 Gut: Nausea, abdominal cramps, vomiting, diarrhea  
 General: Panic, sudden fatigue, chills, feeling of impending doom

#### Examples of potentially life-threatening symptoms that may occur

Throat: Tightening of throat, hoarseness, hacking cough  
 Lung: Shortness of breath, repetitive coughing, wheezing  
 Heart: **THREADY** pulse, passing out, fainting, paleness, blueness

**If reaction is progressing, several of the above body systems may be affected.**

**Note:** While skin symptoms such as itching and hives occur in the majority of food allergic reactions, anaphylaxis does not require the presence of such skin symptoms. **The most**



**dangerous symptoms include breathing difficulties and a drop in blood pressure, causing shock, and are manifested by symptoms like, pallor, dizziness, faintness or passing out.**

**Anaphylaxis is likely when any ONE of the criteria below is fulfilled\*:**

1. Acute onset of an illness (*symptoms may begin within several minutes to two hours after exposure to the allergen*) with involvement of skin/mucosal tissue, such as:

- ✘ Hives,
- ✘ Generalized itch/flush, or
- ✘ Swollen lips/tongue/uvula,

**AND**

Airway compromise, such as:

- ✘ Dyspnea (trouble getting air),
- ✘ Wheeze/bronchospasm, or
- ✘ Stridor (high-pitched breathing noises).

**OR**

Reduced blood pressure or associated symptoms, such as:

- ✘ Hypotonia (decreased muscle tone),
- ✘ Syncope (fainting),
- ✘ Pallor, dizziness, or
- ✘ Blue, weak pulse.

2. Two or more of the following after exposure to suspected or known allergen for that patient (*symptoms may begin within several minutes to two hours after exposure to the allergen*)

- ✘ Skin/mucosal tissue, e.g., hives, generalized itch/flush, swollen lips/tongue/uvula;
- ✘ Airway compromise, e.g., dyspnea (trouble getting air), wheeze/bronchospasm, stridor (high-pitched breathing noises);
- ✘ Associated symptoms, e.g., hypotonia (decreased muscle tone), syncope (faint); or
- ✘ Gastrointestinal symptoms, e.g., crampy abdominal pain, vomiting.

3. Hypotension after exposure to known allergen for that patient (*symptoms may begin within several minutes to two hours after exposure to the allergen*)

- ✘ Infants and children: low systolic BP (age-specific) or >30% drop in systolic BP,\* or
- ✘ Adults: systolic BP, 100 mm Hg or >30% drop from their baseline.

**Note:** Some individuals have an anaphylactic reaction, and the symptoms go away only to return a few hours later. This is called a bi-phasic reaction. Often the symptoms of the bi-phasic reaction occur in the respiratory system and take the individual by surprise. Therefore, according to the American Academy of Allergy, Asthma, and Immunology (AAAAI), after a serious reaction "observation in a hospital setting is necessary for at least four hours after initial symptoms subside because delayed and prolonged reactions may occur even after proper initial treatment." (AAAAI, Position Statement, *Anaphylaxis in schools, and other child-care setting*)

**NOTE: Delegates are not expected to measure and determine a rate of drop in blood pressure as part of epinephrine administration training.**

\* Modified from Symposium on the Definition and Management of Anaphylaxis: Summary report, *JACI*, 2006.

### III. STANDARDS AND PROCEDURES FOR THE EMERGENCY USE OF EPINEPHRINE AUTO-INJECTORS.

An epinephrine auto-injector is a disposable drug delivery device that is easily transportable (about the size of a magic marker) and contains a pre-measured dose of epinephrine. The auto-injector is designed to treat a single anaphylactic episode; and the device must be properly discarded (in compliance with applicable state and federal laws) after its use. It is the responsibility of the parent(s) to provide prescribed epinephrine to the school nurse. As a rule, each student should have two epinephrine auto-injectors available in case subsequent doses of epinephrine are needed to counter a severe reaction.

#### Steps in the Emergency Use of an Epinephrine Auto-Injector:

1. Determine if anaphylaxis is reasonably suspected based on the symptoms identified in the student's IEHP. If there is a reasonable probability that anaphylaxis is occurring or about to occur, then treat the situation like an anaphylactic emergency.
2. Do not leave the student alone. Call 911 and then follow the district's policies and procedures for medical emergencies. Paramedics, the school nurse, the student's parents, and appropriate school administrator(s) should be notified immediately.
3. Most severe allergic reactions in children primarily involve trouble breathing. Children will likely need to sit calmly and upright as they are treated for their breathing problems.

#### **HOWEVER:**

- a) If there is evidence of faintness, loss of consciousness or confusion, lay the child flat. If a child is very ill and needs to be treated in a lying position with legs raised, they should stay in that position, if possible, during transportation to a hospital. Do not change to an upright position, except
  - b) If the child is vomiting, *do not* lay the child flat. If the child is *hypotensive* and vomiting, then he/she should be put down on his/her side, so that the child does not choke on the vomit.
4. Prepare to administer auto-injectable epinephrine, as indicated by the student's health care provider (physician, advanced practice nurse, or physician's assistant) on the IEHP.
    - a) Have the student sit down,
    - b) Reassure the student and avoid moving him or her, and
    - c) Check the auto-injector for expiration date and color. (To be effective, the solution in the auto-injector should be clear and colorless).
  5. Epinephrine Auto-injector Administration Procedure:
    - a) **Grasp the auto-injector in one hand and form a fist around the unit. With the other hand, pull off the safety cap.** (To avoid injecting yourself after removing the cap(s), never place your own fingers or hand over either end of the device. If you accidentally



- inject yourself, then use the back-up auto-injector to treat the student. You should go to the hospital emergency room as well.)
- b) **Hold the tip of the auto-injector near the student's outer thigh.** (The auto-injector can be injected through the student's clothing, if necessary.)
  - c) **Press firmly and hold the tip into the OUTER THIGH so that the auto-injector is perpendicular (at a 90° angle) to the thigh.** You may hear a click.
  - d) **Hold the auto-injector firmly in place for 10-15 seconds.** (After the injection, the student may feel his or her heart pounding. This is a normal reaction.)
  - e) **Remove the auto-injector from the thigh and massage the injection area for several seconds.**
  - f) **Check the tip.** If the needle is exposed, the dose has been delivered. If the needle is not exposed, repeat steps b through e.
  - g) **Dispose of the auto-injector in a "sharps" container or give the expended auto-injector to the paramedics.**
  - h) **Call 911, if not previously called.**
  - i) **Call for the school nurse, if not previously called.**
6. If the anaphylactic reaction is due to an insect sting, remove the stinger (if there is one) as soon as possible after administering the auto-injector. Remove the stinger quickly by scraping with a fingernail, plastic card, or piece of cardboard. Apply an ice pack to the sting area. Do NOT push, pinch, or squeeze, or further imbed the stinger into the skin because such action may cause more venom to be injected into the student.
7. Observe the student. In some circumstances a second administration of epinephrine may be necessary. The school nurse, who is responsible for delegating the administration of epinephrine, must determine that the delegate is competent to administer a second dose of epinephrine in accordance with the written orders from the medical home, the policies and procedures of the district or nonpublic school and the circumstances involved in the emergency.
8. Monitor the student's airway and breathing. If trained in CPR, begin CPR immediately if the student stops breathing.
9. Give a copy of the IEHP to the emergency responders. When emergency responders arrive, tell them the time epinephrine was administered and the dose administered. If the auto-injector has not been disposed of in a sharps container, give the expended auto-injector to the paramedics.

**Note:** Any student who receives epinephrine should be transported to a hospital emergency room, even if symptoms appear to have subsided. If ordered by a health care provider, send a spare auto-injector along with the student to the hospital. A staff member should accompany the child to the hospital and follow procedures in accordance with the district policies regarding the care of students during emergencies.

10. The school nurse should document the incident on the student's health record.

**Notes:**

- ✦ An order for antihistamine administered concurrently with epinephrine does not preclude the emergency administration of epinephrine; however, the antihistamine cannot be given by the delegate even if specified in the IEHP because the statute at N.J.S.A. 18A:40-12.6 only authorizes the delegation of epinephrine.
- ✦ Medical orders that impose a required observation period between administration of an antihistamine and epinephrine are not recommended, in general, and cannot be delegated.

**Storage of Epinephrine Auto-Injectors:**

Epinephrine auto-injectors should be **stored at room temperature** until the marked expiration date, at which time the unit must be replaced. Auto-injectors should not be refrigerated as this could cause the device to malfunction. Auto-injectors should not be exposed to extreme heat (such as in the glove compartment or trunk of a car during the summer) or to direct sunlight. Heat and light shorten the life of the product and can cause the epinephrine to degrade. To be effective, the solution in the auto-injector should be clear and colorless.

P.L. 2007, c. 57 requires that epinephrine auto-injectors be kept in unlocked secure locations to be readily available during an emergency. This will vary depending upon the student's schedule and the circumstances surrounding the possible emergency. In addition to the nurse's office, possible locations for unlocked secure storage of epinephrine include the principal's office, the cafeteria, a classroom, the teacher's room, on the person of the designee, etc. Availability of epinephrine should be addressed in the IEHP) and during the training of the designee by the school nurse.

**REMINDER:** Epinephrine is medication held by the school through the authorization of a health care provider's order. Only school nurses, designees, or students, able to self-medicate, should have epinephrine auto-injectors on their person.



### **III. EMERGENCY FOLLOW-UP**

After epinephrine has been administered, emergency medical care **MUST** be obtained immediately because severely allergic students who have experienced anaphylaxis may need emergency respiratory care, cardiac care, or even resuscitation if they stop breathing. At a minimum, these students will need professional care to determine whether additional epinephrine, steroids, antihistamines, or other treatment is required. Follow-up diagnosis and care by health care professionals after the administration of epinephrine is important for recovery. A delayed or secondary reaction may occur. Therefore, the student needs to remain under medical supervision for at least four hours after an episode of anaphylaxis.

### **IV. POLICIES AND PROCEDURES**

It is strongly recommended but not required that school personnel volunteering to be trained to administer auto-injectable epinephrine also receive instruction and maintain current certification in cardiopulmonary resuscitation from a recognized provider such as the American Red Cross or the American Heart Association.

#### **Statutory Requirements**

N.J.S.A. 18A:40-12.6b and c require district boards of education or chief school administrators of nonpublic schools to implement guidelines and recruit and train volunteers.

N.J.S.A. 18A:40-12.5 requires district boards of education or chief school administrators of nonpublic schools to develop a policy for the emergency administration of epinephrine.

#### **District Board of Education Policies**

N.J.A.C. 6A:16-1.4(a)1 requires district boards of education to develop and adopt written policies, procedures, mechanisms or programs governing the care of any student who becomes injured or ill while at school or during participation in school-sponsored activities.

N.J.A.C. 6A:16-1.4(a)2 requires district boards of education to develop and adopt written policies, procedures, mechanisms or programs governing the transportation and supervision of any student determined to be in need of immediate medical care.

N.J.A.C. 6A:16-1.4(a)7 requires district boards of education to develop and adopt written policies, procedures, mechanisms or programs governing the emergency administration of epinephrine to a student for anaphylaxis pursuant to N.J.S.A. 18A:40-12.5.

N.J.A.C. 6A:16-2.1(a)2iv requires district boards of education to develop and adopt written policies, procedures and mechanisms for the provision of health, safety and medical emergency services and ensure that staff are informed, as appropriate, regarding the administration of medication, including school employees trained and designated by the school nurse to administer epinephrine in an emergency pursuant to N.J.S.A. 18A:40-12.5 and 12.6;

N.J.A.C. 6A:16-2.1(a)4i requires district boards of education to develop and adopt written policies, procedures and mechanisms for the provision of health, safety and medical emergency services and ensure staff are informed as appropriate regarding the provision of health services in emergency situations including the emergency administration of epinephrine pursuant to N.J.S.A. 18A:40-12.5.

N.J.A.C. 6A:7-1.4 requires boards of education shall adopt and implement written educational equity policies that promote equal educational opportunity and foster a learning environment that is free from all forms of prejudice, discrimination and harassment based upon race, creed, color, national origin, ancestry, age, marital status, affectional or sexual orientation, gender, religion, disability or socioeconomic status in the policies, programs and practices of the district board of education.

**Written materials.**

It is the district's responsibility to prepare or obtain these materials and provide them as part of the training.

1. Training Protocols for the Emergency Administration of Epinephrine
2. New Jersey Guidelines for the Care of Students with Food Allergies



## **PART B: RECRUITING DELEGATES FOR THE MANAGEMENT OF ANAPHYLAXIS IN THE SCHOOL SETTING**

New Jersey boards of education or nonpublic school administrators, school nurses, administrators, and staff should work as a team with parents/guardians and the student at risk of anaphylaxis to promote the health, safety, and welfare of students.

New Jersey law requires local boards of education and nonpublic school administrators to develop a policy for the *emergency* administration of epinephrine for anaphylaxis. This policy should be just one of several policies created by the district or nonpublic school regarding the provision of health, safety, and medical emergency services.

The administration of medication in the school setting is the responsibility of the school nurse; however, New Jersey law requires that, as part of emergency planning, additional volunteers be trained to administer epinephrine in the absence of the school nurse. The law also protects the district, the nonpublic school administrator, the school nurse, and the delegate from liability.

### **I. THE ROLE OF SCHOOL BOARDS AND ADMINISTRATORS**

1. Develop requisite policies and procedures outlined in Section IV-Policies and Procedures.
2. Recruitment of delegates requires administrators and school nurses to work collaboratively. There is both an individual and organizational accountability for delegation.
3. Administrators should assist the school nurse in creating circumstances favorable for delegation to occur. These include:
  - a) providing sufficient resources (for example, emergency communication devices and adequate training programs),
  - b) providing opportunity for continuing professional development, including adequate and tangible training for both nurses and staff, and
  - c) creating an environment conducive to teamwork, collaboration, and student-centered care.
4. The NJ Nurse Practice Act authorizes the nurse to delegate and the responsibility accompanying that delegation must follow the law. (N.J.S.A. 45:11-23, and N.J.A.C. 13:37-6.2)

## II. THE ROLE OF THE SCHOOL NURSE

1. The American Nurses Association recognizes that delegation in the school setting is an essential nursing skill. (ANA Position Statement, *Assuring Safe, High Quality Health Care in Pre-K Through 12 Educational Settings*, March 15, 2007)  
<http://www.ana.org/readroom/position/practice/AssuringSafeHealthCarePreK.pdf>
2. New Jersey law specifically recognizes the administration of an epinephrine auto-injector in the school setting as a delegable task by the school nurse. This includes delegation of the auto-injector portion of the Twinject. The nurse must determine that the task, the circumstances, the directions, and supervision are appropriate, and that the risks associated with delegation are minimized. Some considerations for selecting the appropriate delegate include:
  - a) Select a responsible employee who will be reasonably available to the student, particularly where and when anaphylaxis is most likely to occur,
  - b) Consider the knowledge and skills of the individual(s) to whom care may be delegated, and remember that their competency will be enhanced by specific training,
  - c) Consider the proximity and availability of the potential delegate, given the location(s) of the student throughout the school day, the size of the school, and after-hours and off-site school-sponsored functions,
  - d) A student should be assigned as many delegates as necessary to ensure complete back-up coverage, and
  - e) A staff member can serve as a delegate to more than one student, provided he/she has received specific information about each student.
3. A potential delegate is more likely to accept the delegation willingly if the school nurse explains the process with confidence and has approached the candidate thoughtfully. Initial reluctance may be overcome by providing information and reassurance.
4. The school nurse should be proactive in recruiting volunteers that may be appropriate candidates. This means identifying and approaching the candidate, describing the delegation process, and asking, though not compelling, the candidate to accept the delegation.
5. The school nurse provides training to the delegate that includes written instructions and a review of the student's IEHP.
6. The school nurse plans the delegation, based on the student's needs and available resources. The nurse provides periodic and regular evaluation and monitoring of the delegate to ensure an appropriate delegation has been made.
7. School nurses are encouraged to contact the medical home and the school physician to resolve conflicts with written orders that seem to preclude delegation.

8. The school nurse is responsible for documenting the training of each delegate for each student annually, communicating regularly with delegates and updating the Individualized Emergency Healthcare Plan of any student with life-threatening allergies.
9. The law protects the district, the school nurse, and the delegate from liability.



### III. THE ROLE OF THE DELEGATE

The delegate is part of a team whose goal is to provide the best care for students while at school and school-sponsored functions. Although not typically a health care professional, the delegate is critical to the nurse's ability to manage the care of students with severe allergic reactions.

1. Delegates serve as a critical member of the team. A team approach to severe allergy/possible anaphylaxis management is vital i.e., school staff, students, parents, and health care providers should work together to minimize risks and provide a safe educational environment for allergic students.
2. Delegates will participate in training sessions provided by the school nurse that provide clear expectations regarding what to do, what to report, and how to ask for assistance.
3. Delegates will review written instructions from the school nurse and the IEHP including all unique student requirements and characteristics.
4. Delegates will administer epinephrine only in an emergency, when the symptoms described during delegate training are present and the nurse is unavailable.
5. Delegates will ask questions and seek clarification at any time before, during, or after training. Communication is a two-way process and is essential to the provision of services to students during emergencies.
6. Delegates may need to assist students who are permitted to carry and self-administer epinephrine before, during, or after the administration of epinephrine. These students are also entitled to a delegate when unable to self-administer their own epinephrine.
7. The law protects the district, the school nurse, and the delegate from liability.
8. Delegates should be familiar with the following information:
  - a) Allergies can be life-threatening. Exposure to offending allergens can result in *anaphylaxis*. Anaphylaxis, however, is preventable by strict avoidance of offending allergens and treatable by auto-injection of epinephrine (adrenaline) along with medical intervention.
  - b) Epinephrine is most effective for controlling severe allergic reactions in children.
  - c) Through appropriate risk-reduction and allergen-avoidance procedures, the likelihood of the need for epinephrine administration can be minimized.
  - d) Epinephrine is administered in the school setting by auto-injector. An auto-injector is a pre-measured, spring-loaded pen-like device designed for ease of use by non-medical persons in the community. No needle is even visible until the administration is complete.
  - e) The epinephrine auto-injector will be stored in secure and unlocked locations so that it can be readily available for use by the delegate in the event of an emergency. The location of the epinephrine auto-injector will be documented on the IEHP.



## PART C: APPENDIX

### V. RESOURCES

<http://www.foodallergy.org/anaphylaxis/index.html>

<http://www.epipen.com/howtouse.aspx>

<http://www.twinject.com/>

<http://www.redcross.org/services/hss/courses/>

<http://www.aap.org/sections/allergy/child.cfm>

<http://www.aaaai.org/>

**Note:** The American Red Cross has available a training module for administration of auto-injectable epinephrine which can be added to CPR training.

#### **Disclaimer**

*The New Jersey Department of Education does not control or assure the significance, accuracy, or comprehensiveness of the cited resource information. References to resource information in this document are not intended to support any views expressed or products and services offered, nor suggest their importance. The resources identified in the manual are intended to provide schools with links to relevant information for planning, implementing, and evaluating school health procedures. Schools are encouraged to thoroughly assess their needs and investigate programs and materials before adopting them.*

### VI. DEFINITIONS

**Anaphylaxis** - A serious allergic reaction that is rapid in onset and may cause death. (Summary Report of the Second National Institute of Allergy and Infectious Disease/Food Allergy and Anaphylaxis Network symposium on the definition and management of anaphylaxis, *Journal of Allergy and Clinical Immunology*, February 2006).

**Auto-injector** - A pre-measured, spring-loaded pen-like device used to administer epinephrine and designed for ease of use by non-medical persons.

**Certified School Nurse** - A person who holds a current license as a registered professional nurse from the State Board of Nursing and an Educational Services Endorsement, school nurse or school nurse/non-instructional from the Department of Education pursuant to N.J.A.C. 6A:9-13.3 and 13.4.

**Delegation** - Transferring to a competent individual the authority to perform a selected nursing task in a selected situation. The nurse retains accountability for the delegation.

**Delegator** - The person making the decision to delegate the administration of epinephrine.

**Delegate/Designee** - The person receiving designation to administer epinephrine.

**Epinephrine (adrenaline)** - A drug that can be successfully utilized to counteract anaphylaxis.

**Individualized Emergency Healthcare Plan** - A personalized healthcare plan written by the certified school nurse that specifies the delivery of accommodations and services needed by a student in the event of an emergency.

**Individualized Healthcare Plan** - A plan written by the certified school nurse that details accommodations and/or nursing services to be provided to a student because of the student's medical condition based on medical orders written by a health care provider in the student's medical home.

**School-Sponsored Function** - Any activity, event, or program occurring on or off school grounds, whether during or outside of regular school hours, that is organized and/or supported by the school.

973-321-2217  
Edy - Debra  
Dormin  
Rei  
Regi  
X12501

Hadina Dudek  
RN, MA  
CSN  
5000  
x 2398  
973-340-0254



# How to use an EpiPen® (epinephrine injection, USP) Auto-Injector




## 1 PREPARE

Remove the Auto-Injector from the clear carrier tube.  
Flip open the yellow cap of your EpiPen® or the green cap of your EpiPen Jr® carrier tube.  
Tip and slide the auto-injector out of the carrier tube.

Hold the auto-injector in your fist with the orange tip pointing downward.  
Blue to the sky, orange to the thigh®.

With your other hand, remove the blue safety release by pulling straight up without bending or twisting it.



**NEVER-SEE-NEEDLE®** helps with protection. Protects against needle exposure before and after use.

### NOTE:

- The needle comes out of the orange tip.
- To avoid an accidental injection, never put your thumb, fingers or hand over the orange tip. If an accidental injection happens, get medical help right away.

## 2 ADMINISTER

If you are administering to a young child, hold the leg firmly in place while administering an injection.


Place the orange tip against the middle of the outer thigh (upper leg) at a right angle (perpendicular) to the thigh.

Swing and push the auto-injector firmly until it "clicks." The click signals that the injection has started.

Hold firmly in place for 3 seconds (count slowly 1, 2, 3).

Remove the auto-injector from the thigh. The orange tip will extend to cover the needle. If the needle is still visible, do not attempt to reuse it.

Massage the injection area for 10 seconds.



## 3 GET EMERGENCY MEDICAL HELP RIGHT AWAY

You may need further medical attention.

If symptoms continue or recur, you may need to use a second EpiPen® or EpiPen Jr® Auto-Injector.

### INDICATIONS

EpiPen® (epinephrine injection, USP) 0.3 mg or EpiPen Jr® (epinephrine injection, USP) 0.15 mg Auto-Injectors are for the emergency treatment of life-threatening allergic reactions (anaphylaxis) caused by allergens, exercise, or unknown triggers, and for people who are at increased risk for these reactions. EpiPen® or EpiPen Jr® are intended for immediate administration as emergency supportive therapy only. Seek immediate emergency medical help right away.

### IMPORTANT SAFETY INFORMATION

Use EpiPen® or EpiPen Jr® Auto-Injectors right away when you have an allergic emergency (anaphylaxis). **Get emergency medical help right away.** You may need further medical attention. Only a healthcare professional should give additional doses of epinephrine if you need more than two injections for a single anaphylactic episode. EpiPen® or EpiPen Jr® should **only** be injected into the middle of your outer thigh (upper leg), through clothing if necessary. Do not inject into your veins, buttocks, fingers, toes, hands or feet. Hold the leg of young children firmly in place before and during injection to prevent injuries. In case of accidental injection, please seek immediate medical treatment.

**Not actual patient.**


Please see additional Important Safety Information and Indications on the back.  
Please see accompanying Full Prescribing Information and Patient Information.



# Every EpiPen 2-Pak® (epinephrine injection, USP) and Mylan's Authorized Generic For EpiPen Two-Pack comes with an EpiPen® Trainer

Practice with your Trainer repeatedly to become familiar with it.

## The EpiPen® Auto-Injector and Mylan's Authorized Generic For EpiPen®



or

Identify the EpiPen® Trainer by:

### Label

The EpiPen® Trainer is clearly labeled  
TRAINER or TRAINING DEVICE

### Color

The EpiPen® Trainer is shaded grey


EpiPen® and Mylan's  
Authorized Generic  
For EpiPen® are yellow

EpiPen Jr® and Mylan's  
Authorized Generic  
For EpiPen Jr® are green

No Window or Liquid

Reset after each use

## The EpiPen® Trainer



or

The grey EpiPen® Trainer contains no medicine and no needle, and **SHOULD NOT BE USED** during an anaphylactic reaction.

Always have access to two Auto-Injectors in all the places you may need them because some people require a second dose. More than two sequential doses should be administered only under direct medical supervision.

Visit [epipen.com](http://epipen.com) to watch our How to Use video and more.

## IMPORTANT SAFETY INFORMATION (Continued)

Rarely, patients who have used EpiPen® or EpiPen Jr® may develop an infection at the injection site within a few days. Some of these infections can be serious. Call your healthcare professional right away if you have any of the following at an injection site: redness that does not go away, swelling, tenderness, or the area feels warm to the touch.

Tell your healthcare professional about all of your medical conditions, especially if you have asthma, a history of depression, thyroid problems, Parkinson's disease, diabetes, high blood pressure or heart problems, have any other medical conditions, are pregnant or plan to become pregnant, or are breastfeeding or plan to breastfeed. Be sure to also tell your healthcare professional all the medicines you take, especially medicines for asthma. If you have certain medical conditions, or take certain medicines, your condition may get worse or you may have longer lasting side effects when you use EpiPen® or EpiPen Jr®.

Common side effects include fast, irregular or "pounding" heartbeat, sweating, nausea or vomiting, breathing problems, paleness, dizziness, weakness, shakiness, headache, feelings of over excitement, nervousness or anxiety. These side effects usually go away quickly if you lie down and rest. Tell your healthcare professional if you have any side effect that bothers you or that does not go away.

Please see accompanying Full Prescribing Information and Patient Information.

For additional information, please contact us at 300-395-3376.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch) or call 1-800-FDA-1088.

EpiPen®, EpiPen Jr®, EpiPen 2-Pak®, EpiPen Jr 2-Pak®, Now-once-daily, and the Mylan logo are registered trademarks of Mylan Inc. Biotec is a registered trademark of Mylan Specialty L.P. EPI-2016-0079



Better Health  
for a Better World®

The Mylan Better Health for a Better World logo is a registered trademark of Mylan Inc.